

The $\gamma^*\pi^0 \rightarrow \gamma$ form factor

M. A. Ivanov^a and V. E. Lyubovitskij^{a,b}

^a*Bogoliubov Laboratory of Theoretical Physics,
Joint Institute for Nuclear Research, 141980 Dubna, Russia*

^b*Department of Physics, Tomsk State University,
634050 Tomsk, Russia*

Abstract

The $\gamma^*\pi^0 \rightarrow \gamma$ form factor is obtained within the Lagrangian quark model with separable interaction known to provide a good description of the pion observables at low energies. The pion-quarks vertex is chosen in a Gaussian form. The form factor obtained is close to the available experimental data and reaches smoothly the Brodsky-Lepage limit at $Q^2 \approx 10 \text{ GeV}^2$.

Key words: Chiral anomaly; Quark model; Pion transition form factor.

PACS: 12.39.Ki, 13.40.Gp, 14.40.Aq

1 Introduction

The study of neutral pseudoscalar mesons in two-photon reactions provides information on their electromagnetic structure. While the magnitude of $\pi^0 \rightarrow \gamma\gamma$ amplitude is defined by the Adler-Bell-Jackiw axial anomaly [1] which maybe derived from the simple quark diagram with local pion-quark interaction, the knowledge of the form factor with one photon being off mass shell requires the introduction of a nontrivial distribution of quarks inside the pion. The behavior of the off-shell axial anomaly is studied in the $\gamma^*\pi^0 \rightarrow \gamma$ reaction. The data available for this transition form factor in the space-like region $Q^2 < 2.2 \text{ GeV}^2$ are from the CELLO Collab. [2]. New data covering the Q^2 region from 2 GeV^2 up to 20 GeV^2 have been reported by the CLEO Collab. [3]. There is a project to measure this form factor via virtual Compton scattering from a proton target at Jefferson Lab. [4].

The $\gamma^*\pi^0 \rightarrow \gamma$ transition form factor has been studied in the models based on triangle diagrams with nontrivial pion-quark vertex and local quark propagators[5,6], and within a QCD-based model which implements dressing of the vertices and propagators consistent with quark confinement [7]. This transition is a clean test of QCD predictions for exclusive processes. The leading asymptotic behavior $F_{\gamma\pi}(Q^2) \rightarrow 4\pi^2 f_\pi^2/Q^2$ was obtained in [8]. QCD sum rules were applied to calculate this form factor in the region $Q^2 > 1 \text{ GeV}^2$ [9]. A hard scattering approach which includes transverse momentum effects and Sudakov corrections was developed in [10] to describe the large momentum transfer behavior of meson-photon transition form factors.

In this brief note we calculate the $\gamma^*\pi^0 \rightarrow \gamma$ transition form factor within the approach developed in [5]. For the first time, we derive the representation for this form factor from the triangle diagram with dressed pion-quark vertices described by an arbitrary function decreasing rapidly in the Euclidean region. This representation is valid for any momenta of the pion and the photons. We use a Gaussian form for the vertex to make the numerical calculations. The model parameters (the cutoff parameter $\Lambda = 1 \text{ GeV}$ and the constituent quark mass $m_q = 237 \text{ MeV}$) are determined from the π -decay constants and provide a good description of the pion observables at low energies. We find that the form factor obtained is close to the available experimental data and reaches smoothly the Brodsky-Lepage limit at $Q^2 \approx 10 \text{ GeV}^2$.

2 The $\pi^0\gamma\gamma$ vertex function

The basis of the approach [5] is the interaction Lagrangian describing the transition of hadrons into constituent quarks and *vice versa* via an effective vertex function related to the momentum distribution of the constituents. For our purpose, we write the pion Lagrangian as

$$\mathcal{L}_\pi^{\text{int}}(x) = i \frac{g_\pi}{\sqrt{2}} \pi(x) \int dy_1 \int dy_2 f([y_1 - y_2]^2) \delta\left(x - \frac{y_1 + y_2}{2}\right) \bar{q}(y_1) \gamma^5 \tau q(y_2) \quad (1)$$

$$= i \frac{g_\pi}{\sqrt{2}} \pi(x) \int dy f(y^2) \bar{q}(x + y/2) \gamma^5 \tau q(x - y/2).$$

The vertex function $f([y_1 - y_2]^2)$ depending upon the relative coordinate only is chosen to guarantee ultraviolet convergence of the Feynman diagrams. At the same time this function is a phenomenological description of the long-distance QCD interactions between quarks and gluons. The Fourier transform of the vertex function in Minkowsky space is defined as $f(y^2) = \int d^4k / (2\pi)^4 f(k^2)$. The gauging of the nonlocal Lagrangian Eq. (1) has been done in [5] and, in more general form, in [11] by using the time-ordering P -exponential. It leads to the appearance of additional diagrams in some electromagnetic processes. However, they do not appear in the anomalous processes like $\pi^0 \rightarrow \gamma\gamma$ or $\gamma \rightarrow 3\pi$.

The coupling constant g_π is determined from the so-called *compositeness condition* or, equivalently, from the normalization of the electromagnetic form factor to one at the origin. Its value as well the value of pion weak coupling constant are defined by

$$\left(\frac{3g_\pi^2}{4\pi^2} \right)^{-1} \simeq \frac{1}{4} \int_0^\infty du u f^2(-u) \frac{(3m_q^2 + 2u)}{(m_q^2 + u)^3}. \quad (2)$$

$$f_\pi \simeq \frac{3g_\pi}{4\pi^2} m_q \int_0^\infty du u f(-u) \frac{1}{(m_q^2 + u)^2}.$$

Here, the pion mass is neglected since its magnitude is much less than the cut-off parameter (see, [5]).

The form factor which defines the two-photon decay of the pion with the pion and photon momenta being off mass shell is written in the form

$$G_{\pi\gamma\gamma}(p^2, q_1^2, q_2^2) = m_q \cdot \frac{g_\pi}{2\sqrt{2}\pi^2} R_{\gamma\pi}(p^2, q_1^2, q_2^2) \quad (3)$$

$$R_{\gamma\pi}(p^2, q_1^2, q_2^2) = \int \frac{d^4k}{\pi^2 i} \frac{f(k^2)}{D_0} \quad (4)$$

$D_0 = [m_q^2 - (k + p/2)^2][m_q^2 - (k - p/2)^2][m_q^2 - (k + (q_1 - q_2)/2)^2]$. Using the Feynman α -parametrization for $1/D_0$ one finds

$$R_{\gamma\pi}(p^2, q_1^2, q_2^2) = 2 \int d^3\alpha \delta \left(1 - \sum_{i=1}^3 \alpha_i \right) \int \frac{d^4k}{\pi^2 i} \frac{f(k^2)}{\left[m_q^2 - D_1 - (k + r)^2 \right]^3} \quad (5)$$

where $D_1 = \alpha_1 \alpha_2 p^2 + \alpha_1 \alpha_3 q_1^2 + \alpha_2 \alpha_3 q_2^2$ and $r = q_2(1 - 2\alpha_2)/2 - q_1(1 - 2\alpha_1)/2$. Then we use the Cauchy theorem for the function

$$f(k^2) = \oint \frac{dz}{2\pi i} \frac{f(z)}{z - k^2},$$

and, again, the Feynman α -parametrization. Performing the integration over k gives

$$R_{\gamma\pi}(p^2, q_1^2, q_2^2) = \int_0^\infty dt \left(\frac{t}{1+t} \right)^2 \int d^3\alpha \delta \left(1 - \sum_{i=1}^3 \alpha_i \right) \left\{ -f' \left[-t(m_q^2 - D_1) + \frac{t}{1+t} r^2 \right] \right\}$$

where $r^2 = -D_1 + p^2(1 - 2\alpha_3)/4 + (q_1^2 + q_2^2)\alpha_3/2$. Note that this representation is valid for any function $f(k^2)$ and convenient for numerical calculations.

The two-photon decay coupling constant is obtained from Eq. (3) for both q_1^2 and q_2^2 being equal to zero

$$g_{\pi\gamma\gamma} = G_{\pi\gamma\gamma}(m_\pi^2, 0, 0) \simeq m_q \frac{g_\pi}{2\sqrt{2}\pi^2} \int_0^\infty du u f(-u) \frac{1}{(m_q^2 + u)^3}. \quad (6)$$

If we use the Gaussian form for the vertex function $f(k^2) = \exp\{k^2/\Lambda^2\}$ the best fit is found to be $f_\pi = 132$ MeV ($f_\pi^{\text{expt}} = 132$ MeV) and $g_{\pi\gamma\gamma} = 0.261$ GeV⁻¹ ($g_{\pi\gamma\gamma}^{\text{expt}} = 0.276$ GeV⁻¹) with the adjustable parameters being equal to $\Lambda = 1$ GeV and $m_q = 237$ MeV.

3 The transition form factor

Defining the transition form factor by

$$F_{\gamma\pi}(Q^2) = e^2 G_{\pi\gamma\gamma}(m_\pi^2, -Q^2, 0) \simeq e^2 \frac{g_\pi}{2\sqrt{2}\pi^2} \frac{m_q}{\Lambda^2} R_{\gamma\pi}(Q^2) \quad (7)$$

Then the decay width is written as

$$\Gamma(\pi^0 \rightarrow \gamma\gamma) = \frac{m_\pi^3}{64\pi} F_{\gamma\pi}^2(0). \quad (8)$$

Our results for $(m_\pi^3/64\pi)F_{\gamma\pi}^2(Q^2)$ are shown in Fig.1 comparing with the experimental data and the monopole fit

$$F_{\gamma\pi}^{\text{fit}}(Q^2) = \frac{e^2 g_{\pi\gamma\gamma}}{1 + Q^2/\Lambda_\pi^2} \quad \Lambda_\pi = 0.6 \text{ GeV}. \quad (9)$$

One can see that the form factor is close to the available experimental data.

The radius for $\gamma^*\pi^0 \rightarrow \gamma$ transition is defined by

$$\langle r_{\gamma\pi}^2 \rangle = -6 \frac{F'_{\gamma\pi}(0)}{F_{\gamma\pi}(0)} \quad (10)$$

where

$$F_{\gamma\pi}(0) = \int_0^\infty du u \frac{f(-u)}{(m_q^2 + u)^3} \quad F'_{\gamma\pi}(0) = -\frac{m_q^2}{2} \int_0^\infty du u \frac{f(-u)}{(m_q^2 + u)^5}. \quad (11)$$

With this definition one finds $r_{\gamma\pi}=0.65$ fm which is the experimental value having an error of 0.03 fm.

We compare our result for the function $Q^2 F_{\gamma\pi}(Q^2)/F_{\gamma\pi}(0)$ with the Brodsky-Lepage limit $4\pi^2 f_\pi^2 \approx 0.68 \text{ GeV}^2$ in the Fig.2. One can see that this function reaches smoothly the Brodsky-Lepage limit at $Q^2 \approx 10 \text{ GeV}^2$.

Summarizing we have calculated the pion weak decay constant, the two-photon decay width, as well as the form factor of the $\gamma^*\pi^0 \rightarrow \gamma$ -transition. The two adjustable parameters, the range parameter Λ appearing in the Gaussian, and

the constituent quark mass m_q , have been fixed by fitting the experimental data for the pion decay constants. We find that the $\gamma\pi$ - form factor is close to the available experimental data and smoothly reaches the Brodsky-Lepage limit at $Q^2 \approx 10 \text{ GeV}^2$.

4 Acknowledgements

We acknowledge helpful discussions with M.P. Locher during our stay at PSI. This work was supported in part by the INTAS Grant 94-739 and the Russian Fund of Fundamental Research under contract 96-02-17435-a.

References

- [1] S. Adler, Phys. Rev. **177** (1969) 2426;
J. Bell and R. Jackiw, Nuovo Cimento **A60** (1969) 47.
- [2] CELLO Collaboration, H.J. Behrend et al., Z. Phys. **C49** (1991) 401.
- [3] V. Savinov (CLEO Collab.), hep-ex/9507005, 1995.
- [4] A. Afanasev, Preprint JLAB-THY-96-01, hep-ph/9608305, 1996.
- [5] I.V. Anikin, M.A. Ivanov, N.B. Kulimanova, V.E. Lyubovitskij,
Z. Phys. **C65** (1995) 681.
- [6] H. Ito, W.W. Buck, F. Gross, Phys. Lett. **B287** (1992) 23.
- [7] M.R. Frank, K.L. Mitchell, C.D. Roberts, P.C. Tandy,
Phys. Lett. **B359** (1995) 17.
- [8] S.J. Brodsky and G.P. Lepage, Phys. Rev. **D24** (1981) 1808.
- [9] A.V. Radyushkin and R. Ruskov, Phys. Lett. **B374** (1996) 173.
- [10] P. Kroll and M. Raulf, Phys. Lett. **B397** (1996) 848;
R. Jakob, P. Kroll, M. Raulfs, J. Phys. **G22** (1996) 45.
- [11] M.A. Ivanov, M.P. Locher, and V.E. Lyubovitsij,
Few-Body Systems, **21** (1996) 131.

List of figures

Fig.1. Our calculated form of $m_\pi^3 F_{\gamma\pi}^2(Q^2)/64\pi$ (dashed line) compared with experimental data and their fit (solid line) [2,3].

Fig.2. Our calculated form of $Q^2 F_{\gamma\pi}(Q^2)/F_{\gamma\pi}(0)$ (dashed line) compared with experimental fit (solid line) and the Brodsky-Lepage limit ($\approx 0.68 \text{ GeV}^2$) [8].



